




Preliminary Examination Paper 1 Solutions

Qn	Solution	Remarks
<p>1</p>	$\frac{2w-5}{w^2-3} > 0$  <p style="text-align: center;">OR</p>  $-\sqrt{3} < w < \sqrt{3} \quad \text{or} \quad w > \frac{5}{2}$ $\frac{(2 y -5)\sin x}{y^2-3} \leq 0,$ <p>Since $\sin x < 0$ for $\pi < x \leq \frac{3\pi}{2}$, $\frac{2 y -5}{y^2-3} \geq 0$</p> <p>From above, $-\sqrt{3} < y < \sqrt{3}$ or $y \geq \frac{5}{2}$</p> $0 \leq y < \sqrt{3} \quad \text{or} \quad y \geq \frac{5}{2}$ $-\sqrt{3} < y < \sqrt{3} \quad \text{or} \quad y \geq \frac{5}{2} \quad \text{or} \quad y \leq -\frac{5}{2}$	
<p>2</p>	 <p>Method 1 - Integration</p> <p>Volume of solid S</p> $= \pi \int_0^e \left(\frac{x}{e}\right)^2 dx - \pi \int_1^e (\ln x)^2 dx$ $= \frac{\pi}{e^2} \int_0^e x^2 dx - \pi \int_1^e (\ln x)^2 dx$ $= \frac{\pi}{e^2} \left[\frac{x^3}{3} \right]_0^e - \pi \int_1^e (\ln x)^2 dx$ <p>Method 2 - Volume of Cone</p> <p>Volume of solid S</p> $= \text{Vol of cone} - \pi \int_1^e (\ln x)^2 dx$ $= \frac{1}{3} \pi (1)^2 (e) - \pi \int_1^e (\ln x)^2 dx$ $= \frac{1}{3} \pi e - \pi \int_1^e (\ln x)^2 dx$	

	$= \frac{\pi}{e^2} \left[\frac{e^3}{3} \right] - \pi \int_1^e (\ln x)^2 dx$ $= \frac{1}{3} \pi e - \pi \int_1^e (\ln x)^2 dx$ $\int_1^e (\ln x)^2 dx$ $= \left[x(\ln x)^2 \right]_1^e - 2 \int_1^e \ln x dx \quad u_1 = (\ln x)^2 \quad \frac{dv_1}{dx} = 1$ $= \left[x(\ln x)^2 \right]_1^e - 2 \left([x \ln x]_1^e - \int_1^e 1 dx \right) \quad \frac{du_1}{dx} = 2(\ln x) \left(\frac{1}{x} \right) \quad v_1 = x$ $= \left[x(\ln x)^2 \right]_1^e - 2[x \ln x]_1^e + 2 \int_1^e 1 dx \quad u_2 = \ln x \quad \frac{dv_2}{dx} = 1$ $= \left[x(\ln x)^2 \right]_1^e - 2[e \ln e - \ln 1]_1^e + 2[x]_1^e \quad \frac{du_2}{dx} = \frac{1}{x} \quad v_2 = x$ $= \left[x(\ln x)^2 \right]_1^e - 2[e \ln e - \ln 1] + 2[e - 1]$ $= e - 2e + 2e - 2$ $= e - 2$ <p>Hence, volume of solid S</p> $= \frac{1}{3} \pi e - \pi(e - 2)$ $= \frac{1}{3} \pi e - \pi e + 2\pi$ $= 2\pi - \frac{2}{3} \pi e$ $= \frac{2}{3} \pi(3 - e)$	
3 (i)	$2\,500\,000\,000 = 1000\,000(2)^{n-1}$ $2^{n-1} = 2500$ $n-1 = \frac{\ln 2500}{\ln 2} = 11.2877$ $n = 12.2877$ <p>His net worth will first exceed 2.5 billion when $n = 13$ The year $1993 + (13 - 1)(1) = 2005$ or $1993 + 13 - 1 = 2005$</p>	
3 (ii)	$100\,000 + 1000 + 1000(1.5) + 1000(1.5^2) + \dots + (15 \text{ terms}) =$ $100\,000 + \frac{1000(1.5^{15} - 1)}{1.5 - 1} = \$973\,787.7808 = \$973\,788$	
4 (a)	$g(x) = \frac{1}{\left(\frac{1}{\sqrt{2}} \cos x - \frac{1}{\sqrt{2}} \sin x \right) \left(\frac{1}{\sqrt{2}} \cos x + \frac{1}{\sqrt{2}} \sin x \right)} = \frac{1}{\left(\frac{1}{2} \cos^2 x - \frac{1}{2} \sin^2 x \right)} =$	

	$\frac{1}{\frac{1}{2}(\cos^2 x - \sin^2 x)} = \frac{2}{\cos 2x} \approx \frac{2}{(1 - \frac{(2x)^2}{2})} = \frac{2}{1 - 2x^2} = 2(1 - 2x^2)^{-1}$ $\approx 2(1 + 2x^2) = 2 + 4x^2$ <p>m must be sufficiently small for $g(x) \approx 2 + ax + bx^2$</p>	
4 (b)	$(1 - x^2)^{-1/2} = 1 + \frac{(-\frac{1}{2})}{1}(-x^2) + \frac{(-\frac{1}{2})(-\frac{3}{2})}{2!}(-x^2)^2 +$ $\frac{(-\frac{1}{2})(-\frac{3}{2})(-\frac{5}{2})}{3!}(-x^2)^3 = 1 + \frac{1}{2}x^2 + \frac{3}{8}x^4 + \frac{5}{16}x^6 + \dots$ $\cos^{-1} x = \int \frac{-1}{\sqrt{1-x^2}} dx = -\int (1 + \frac{1}{2}x^2 + \frac{3}{8}x^4 + \frac{5}{16}x^6 + \dots) dx$ $\approx -(x + \frac{1}{6}x^3 + \frac{3}{40}x^5 + \frac{5}{112}x^7) + C$ <p>When $x = 0$, $\cos^{-1} 0 = \frac{\pi}{2} = C$</p> $\cos^{-1}(x) = -(x + \frac{1}{6}x^3 + \frac{3}{40}x^5) + \frac{\pi}{2}$	
5	$\frac{1}{3^r} \left(\frac{u_{r+1}}{3} - u_r \right) = 2r$ $\sum_{r=1}^n \frac{u_{r+1}}{3^{r+1}} - \frac{u_r}{3^r} = 2 \sum_{r=1}^n r$ $\frac{u_2}{3^2} - \frac{u_1}{3^1}$ $+ \frac{u_3}{3^3} - \frac{u_2}{3^2}$ $+ \dots$ $+ \frac{u_n}{3^n} - \frac{u_{n-1}}{3^{n-1}}$ $+ \frac{u_{n+1}}{3^{n+1}} - \frac{u_n}{3^n} = 2 \left(\frac{n}{2} (1+n) \right)$ $\frac{u_{n+1}}{3^{n+1}} - \frac{1}{3} = n(n+1)$ $u_{n+1} = 3^{n+1} \left(n(n+1) + \frac{1}{3} \right) = 3^n (3n^2 + 3n + 1)$	
6	<p>Rate of change = rate of growth – rate of decrease Rate of change = rate of earning interest – rate of withdrawal</p> $\frac{dM}{dt} = 0.05M - 4000$	

$$\int \frac{1}{0.05M - 4000} dM = \int 1 dt$$

$$\frac{1}{0.05} \ln |0.05M - 4000| = t + C$$

$$\ln |0.05M - 4000| = \frac{t}{20} + \frac{C}{20}$$

$$|0.05M - 4000| = e^{\frac{t}{20} + \frac{C}{20}}$$

$$0.05M - 4000 = Ae^{\frac{t}{20}} \text{ where } A = \pm e^{\frac{C}{20}}$$


$$M = 80000 + 20Ae^{\frac{t}{20}}$$

$$\text{When } t = 0, M = K$$

$$K = 80000 + 20A$$

$$20A = K - 80000$$

$$\text{Hence } M = 80000 + (K - 80000)e^{\frac{t}{20}}$$



Money is completely withdrawn if $K < 80000$

7(i) Let $x = 5 \sin \theta$

$$\frac{dx}{d\theta} = 5 \cos \theta$$

$$\int \sqrt{25 - x^2} dx$$

$$= \int \sqrt{25 - (5 \sin \theta)^2} (5 \cos \theta) d\theta$$

$$= \int \sqrt{25 - 25 \sin^2 \theta} (5 \cos \theta) d\theta$$

$$= \int \sqrt{25(1 - \sin^2 \theta)} (5 \cos \theta) d\theta$$

$$= \int \sqrt{25(\cos^2 \theta)} (5 \cos \theta) d\theta$$

$$= \int (5 \cos \theta)^2 d\theta$$


$$= 25 \int \cos^2 \theta d\theta$$

$$= \frac{25}{2} \int (1 + \cos 2\theta) d\theta$$

$$= \frac{25}{2} \left(\theta + \frac{1}{2} \sin 2\theta \right) + c$$

$$= \frac{25}{2} (\theta + \sin \theta \cos \theta) + c$$

$$= \frac{25}{2} \left(\sin^{-1} \frac{x}{5} + \left(\frac{x}{5} \right) \left(\frac{\sqrt{25 - x^2}}{5} \right) \right) + c$$



$$\text{Let } x = 5 \sin \theta$$

$$\sin \theta = \frac{x}{5}$$

$$\theta = \sin^{-1} \frac{x}{5}$$

$$\cos \theta = \frac{1}{5} \sqrt{25 - x^2}$$

	$= \frac{25}{2} \sin^{-1} \frac{x}{5} + \frac{1}{2} x \sqrt{25-x^2} + c$	
7(ii)	$x^2 + (y-b)^2 = 25$ $(y-b)^2 = 25 - x^2$ <p>Since $y < 0$, $y - b < 0$,</p> $y - b = -\sqrt{25 - x^2}$ $y = b - \sqrt{25 - x^2}$ <p>Area of region R</p> $= \left \int_0^a b - \sqrt{25 - x^2} dx \right $ $= -\int_0^a b - \sqrt{25 - x^2} dx$ $= \int_0^a \sqrt{25 - x^2} dx - \int_0^a b dx$ $= \left[\frac{25}{2} \sin^{-1} \frac{x}{5} + \frac{1}{2} x \sqrt{25 - x^2} \right]_0^a - [bx]_0^a$ $= \frac{25}{2} \sin^{-1} \frac{a}{5} + \frac{1}{2} a \sqrt{25 - a^2} - ab$ $= \frac{1}{2} a \sqrt{25 - a^2} + \frac{25}{2} \sin^{-1} \frac{a}{5} - a \sqrt{25 - a^2}$ $= \frac{25}{2} \sin^{-1} \frac{a}{5} - \frac{1}{2} a \sqrt{25 - a^2}.$	<p>Substituting $x = a$ and $y = 0$ into the equation</p> $x^2 + (y-b)^2 = 25,$ <p>we have</p> $a^2 + (0-b)^2 = 25$ $b = \sqrt{25 - a^2}$
8 (i)	$z = k + i$ $z^2 = (k + i)^2 = k^2 + 2(k)(i) + (i)^2 = (k^2 - 1) + (2k)i$ $z^3 = (k + i)^3 = k^3 + 3(k)^2(i) + 3(k)(i)^2 + (i)^3$ $= (k^3 - 3k) + (3k^2 - 1)i$ $z^3 - iz^2 - 2z - 4i = 0$ $[(k^3 - 3k) + (3k^2 - 1)i] - i[(k^2 - 1) + (2k)i] - 2[k + i] - 4i = 0$ $[(k^3 - 3k) + 2k - 2k] + i[(3k^2 - 1) - (k^2 - 1) - 2 - 4] = 0$ $(k^3 - 3k) + i(2k^2 - 6) = 0$ $k(k^2 - 3) = 0 \text{ and } 2k^2 - 6 = 0$ $(k = 0 \text{ or } k = \pm\sqrt{3}) \text{ and } k = \pm\sqrt{3}$ <p>Hence, $k = \pm\sqrt{3}$</p>	
8 (ii)	$z = \sqrt{3} + i \quad (\because k > 0)$ $ z = \sqrt{1+3} = 2$ $\arg(z) = \frac{\pi}{6}$	

Method 1: By Polar Form & Trigonometry

$$z = 2e^{i\pi/6} = 2\left(\cos\frac{\pi}{6} + i\sin\frac{\pi}{6}\right)$$

$$z^n = 2^n e^{in\pi/6} = 2^n\left(\cos\frac{n\pi}{6} + i\sin\frac{n\pi}{6}\right)$$

$$z^n \text{ is real} \Leftrightarrow \sin\frac{n\pi}{6} = 0$$

$$\Leftrightarrow \frac{n\pi}{6} = k\pi, \text{ where } k \in \mathbb{Z}$$

$$\Leftrightarrow n = 6k, \text{ where } k \in \mathbb{Z}$$

Hence, $n = 0, \pm 6, \pm 12, \pm 18, \dots$

Method 2: By Properties of $\arg(z)$

$$\arg(z^n) = n \arg(z) = \frac{n\pi}{6}$$

z^n is real, the point representing z^n on the Argand diagram is on the x -axis.

$$\text{Thus, } \arg(z^n) = \frac{n\pi}{6} = k\pi, \text{ where } k \in \mathbb{Z}$$

$$\therefore n = 6k, \text{ where } k \in \mathbb{Z}$$

i.e. $n = 0, \pm 6, \pm 12, \pm 18, \dots$

Given $|z^n| > 100$.

$$|z^n| = |z|^n = 2^n$$

Hence, $2^n > 100$

But n is a multiple of 6. We then have

$$2^6 = 64 < 100$$

$$2^{12} = 4096 > 100$$

The least value of n is then 12.

9

Let P_n be the statement $\sum_{r=1}^n \frac{3r+1}{r(r+1)(r+2)} = \frac{n(7n+9)}{4(n+1)(n+2)}$ for

all integers $n \geq 1$.

$$\text{When } n = 1, \quad \text{LHS} = \sum_{r=1}^1 \frac{3r+1}{r(r+1)(r+2)} = \frac{3+1}{1(2)(3)} = \frac{2}{3}$$

$$\text{RHS} = \frac{7+9}{4(2)(3)} = \frac{2}{3} = \text{LHS}$$

$\therefore P_1$ is true.

Assume that P_k is true for **some** positive integer $k, k \geq 1$,

$$\text{i.e. } \sum_{r=1}^k \frac{3r+1}{r(r+1)(r+2)} = \frac{k(7k+9)}{4(k+1)(k+2)}$$

Need to prove P_{k+1} is true,

i.e.
$$\sum_{r=1}^{k+1} \frac{3r+1}{r(r+1)(r+2)} = \frac{(k+1)(7k+16)}{4(k+2)(k+3)}.$$

LHS of $P_{k+1} = \sum_{r=1}^{k+1} \frac{3r+1}{r(r+1)(r+2)}$

$$= \sum_{r=1}^k \frac{3r+1}{r(r+1)(r+2)} + \frac{3(k+1)+1}{(k+1)(k+2)(k+3)}$$

$$= \frac{k(7k+9)}{4(k+1)(k+2)} + \frac{3k+4}{(k+1)(k+2)(k+3)}$$

$$= \frac{7k^3 + 30k^2 + 39k + 16}{4(k+1)(k+2)(k+3)}$$

$$= \frac{(7k^2 + 23k + 16)(k+1)}{4(k+1)(k+2)(k+3)}$$

$$= \frac{(k+1)(7k+16)}{4(k+2)(k+3)} = \text{RHS}$$

$\therefore P_k$ is true $\Rightarrow P_{k+1}$ is true.

Since P_1 is true, and P_k is true $\Rightarrow P_{k+1}$ is true, by the Principle of Mathematical Induction, $\sum_{r=1}^n \frac{3r+1}{r(r+1)(r+2)} = \frac{n(7n+9)}{4(n+1)(n+2)}$ is true for all integers $n \geq 1$.

9 (i)
$$\frac{n(7n+9)}{4(n+1)(n+2)} = \frac{7}{4} - \frac{6n+7}{2(n+1)(n+2)}$$

Since n is a positive integer, $\frac{6n+7}{2(n+1)(n+2)} > 0$


$$\therefore \sum_{r=1}^n \frac{3r+1}{r(r+1)(r+2)} = \frac{n(7n+9)}{4(n+1)(n+2)} = \frac{7}{4} - \frac{6n+7}{2(n+1)(n+2)} < \frac{7}{4}$$

Alternatively,

$$\sum_{r=1}^n \frac{3r+1}{r(r+1)(r+2)} = \sum_{r=1}^n \frac{1}{2r} + \frac{2}{r+1} - \frac{5}{2(r+2)} = \dots = \frac{7}{4} - \frac{1}{2(n+1)} - \frac{5}{2(r+2)} < \frac{7}{4}$$

Since n is a positive integer, $-\frac{1}{2(n+1)} - \frac{5}{2(n+2)} < 0$

<p>9 (ii)</p>	$(r+1)^3 = r^3 + 3r^2 + 3r + 1$ $r(r+1)(r+2) = r^3 + 3r^2 + 2r$ $\therefore (r+1)^3 > r(r+1)(r+2)$ $\sum_{r=1}^n \frac{3r}{(r+1)^3} < \sum_{r=1}^n \frac{3r}{r(r+1)(r+2)} < \sum_{r=1}^n \frac{3r+1}{r(r+1)(r+2)} < \frac{7}{4}.$	
<p>10</p>	$y = \frac{x^2 - 4k^2}{x - k} \quad \text{where } k \text{ is a constant such that } k \neq 0$ $xy - ky = x^2 - 4k^2$ $x^2 - xy + (ky - 4k^2) = 0$ <p>x is real \Rightarrow discriminant ≥ 0</p> $y^2 - 4(ky - 4k^2) \geq 0$ $y^2 - 4ky + 16k^2 \geq 0$ $(y - 2k)^2 + 12k^2 \geq 0$ <p>This inequality is true for all values of y. Therefore y can take the set of all real numbers.</p> <p><i>Alternative Method:</i></p> $\frac{dy}{dx} = 0$ $\frac{x^2 - 2xk + 4k^2}{(x - k)^2} = 0$ $x^2 - 2xk + 4k^2 = 0$ <p>discriminant = $-12k^2 < 0$ no real roots. Hence, no turning pts. $y \in \mathbb{R}$.</p>	

<p>10 (i)</p>	$y = \frac{x^2 - 4k^2}{x - k} = x + k - \frac{3k^2}{x - k}$ <p>Asymptotes: $y = x + k$ and $x = k$</p> <p>Points of intercept with axes: $(0, 4k)$, $(-2k, 0)$, $(2k, 0)$</p> 	
<p>10 (ii)</p>	$\int_{-1}^1 f(x) dx = 2 \int_0^1 f(x) dx = 2 \int_0^1 \left(x + k - \frac{3k^2}{x - k} \right) dx$ $= 2 \left[\frac{x^2}{2} + kx - 3k^2 \ln x - k \right]_0^1$ $= 2 \left[\frac{1}{2} + k - 3k^2 \ln 1 - k + 3k^2 \ln -k \right]$ $= 1 + 2k + 6k^2 \ln \frac{k}{k - 1}.$	
<p>10 (iii)</p>	$y = \frac{x^2 - 4k^2}{x - k} \rightarrow y = \frac{\left(\frac{x}{2}\right)^2 - 4k^2}{\left(\frac{x}{2}\right) - k} \rightarrow y = \frac{x^2 - 16k^2}{2x - 4k}$ $\rightarrow y = \frac{(x + 2k)^2 - 16k^2}{2(x + 2k) - 4k} = \frac{(x + 2k)^2 - 16k^2}{2x}$	


11 (i)	$\mathbf{a} \times \mathbf{b} = 4\mathbf{a} \times \mathbf{c}$ $(\mathbf{a} \times \mathbf{b}) - (4\mathbf{a} \times \mathbf{c}) = 0$ $(\mathbf{a} \times \mathbf{b}) - (\mathbf{a} \times 4\mathbf{c}) = 0$ $\mathbf{a} \times (\mathbf{b} - 4\mathbf{c}) = 0$ \mathbf{a} is parallel to $\mathbf{b} - 4\mathbf{c}$ $\mathbf{b} - 4\mathbf{c} = \alpha \mathbf{a}$	
(ii)	$\frac{1}{2} \mathbf{a} \times \mathbf{b} = \sqrt{126}$ $\frac{1}{2} 4\mathbf{a} \times \mathbf{c} = \sqrt{126}$ $ \mathbf{a} \times \mathbf{c} = \frac{\sqrt{126}}{2}$ $\left \left(\frac{\mathbf{b} - 4\mathbf{c}}{\sqrt{3}} \right) \times \mathbf{c} \right = \frac{\sqrt{126}}{2}$ $ (\mathbf{b} \times \mathbf{c}) - (4\mathbf{c} \times \mathbf{c}) = \frac{\sqrt{3}\sqrt{126}}{2}$ $ (\mathbf{b} \times \mathbf{c}) = \frac{\sqrt{378}}{2}$ <p>Alternatively,</p> $ \mathbf{b} \times \mathbf{c} = \left \mathbf{b} \times \left(\frac{\mathbf{b} - \sqrt{3}\mathbf{a}}{4} \right) \right = \frac{1}{4} \mathbf{b} \times \mathbf{b} - \mathbf{b} \times \sqrt{3}\mathbf{a} $ $= \frac{\sqrt{3}}{4} \mathbf{b} \times \mathbf{a} = \frac{\sqrt{3}}{2} \cdot \frac{1}{2} \mathbf{b} \times \mathbf{a} = \frac{\sqrt{3}}{2} \cdot \sqrt{126}$ $\therefore (\mathbf{b} \times \mathbf{c}) = \frac{\sqrt{378}}{2}$	
(iii)	Area of parallelogram with adjacent sides OB and OC .	
(iv)	$(\mathbf{b} - 4\mathbf{c}) \cdot (\mathbf{b} - 4\mathbf{c}) = 3 \mathbf{a} ^2$ $ \mathbf{b} ^2 - 8\mathbf{b} \cdot \mathbf{c} + 16 \mathbf{c} ^2 = 3 \mathbf{a} ^2$ $\mathbf{b} \cdot \mathbf{c} = -\frac{10}{8}$ $\cos \theta = \frac{\mathbf{b} \cdot \mathbf{c}}{ \mathbf{b} \mathbf{c} } = \frac{-\frac{10}{8}}{1(2)}$ $\theta = 128.7^\circ$	
12 (i)	$z^5 - i = 0$ $z^5 = i$ $z^5 = e^{i\pi/2} = e^{i(2k\pi + \frac{\pi}{2})}$, where $k \in \mathbb{Z}$ $z = e^{i(\frac{2k\pi}{5} + \frac{\pi}{10})}$ Putting $n = -2, -1, 0, 1, 2$	

$$z = e^{-\frac{7\pi i}{10}}, e^{-\frac{3\pi i}{10}}, e^{\frac{\pi i}{10}}, e^{\frac{\pi i}{2}}, e^{\frac{9\pi i}{10}}$$

Given $-\pi < \arg(z_1) < \arg(z_2) < \arg(z_3) < \arg(z_4) < \arg(z_5) \leq \pi$.

i.e. $z_1 = e^{-\frac{7\pi i}{10}}, z_2 = e^{-\frac{3\pi i}{10}}, z_3 = e^{\frac{\pi i}{10}}, z_4 = e^{\frac{\pi i}{2}}, z_5 = e^{\frac{9\pi i}{10}}$

Let the points P_1, P_2, P_3, P_4 and P_5 on the Argand diagram represents the complex numbers $z_1, z_2, z_3, z_4,$ and z_5 .



The locus $|z - z_2| = |z - z_3|$ is a perpendicular bisector of the line segment P_2P_3 where $P_2 \equiv z_2$ and $P_3 \equiv z_3$.

12
(ii)

Since OP_2P_3 is an isosceles triangle, the perpendicular cuts through the origin and bisect the angle P_2OP_3 .


Thus the perpendicular bisector is inclined at angle $\frac{\pi}{10}$ radian

below the positive x -axis.

Hence, the Cartesian equation of the locus $|z - z_2| = |z - z_3|$ is

$$y = -x \tan\left(\frac{\pi}{10}\right) \text{ OR } y = x \tan\left(\frac{9\pi}{10}\right)$$

$$\arg(z - z_1) = \arg(z_4)$$




$$z \equiv \overline{OP}$$

$$z_1 \equiv \overline{OP_1}, \quad z - z_1 \equiv \overline{OP} - \overline{OP_1} \text{ i.e. } z - z_1 \equiv \overline{P_1P}$$

The least value of $|z - z_1|$ is the shortest distance from P_1 to the perpendicular bisector (line OP_5).

$$\frac{P_1N}{OP_1} = \sin \frac{4\pi}{10}$$

$$\begin{aligned} P_1N &= OP_1 \sin \frac{4\pi}{10} \\ &= \sin \frac{2\pi}{5} = 0.951 \end{aligned}$$



12 (iii) The locus $\arg(z - z_1) = \arg(z_4)$ is a half-line with its initial point at P_1 and above and excluding P_1 , parallel to the y -axis. (Refer to diagram.)

12 (iii) The intersection point between the 2 loci has the same x -coordinates as the point P_1 , i.e. $x = \cos \frac{-7\pi}{10} = -\cos \frac{3\pi}{10}$.



Substituting $x = -\cos \frac{3\pi}{10}$ into the equation of the perpendicular

bisector $y = -x \tan(\frac{\pi}{10})$, we have

$$y = -(-\cos \frac{3\pi}{10}) \tan \frac{\pi}{10} = \cos \frac{3\pi}{10} \tan \frac{\pi}{10} = 0.191$$

$$x = -\cos \frac{3\pi}{10} = -0.588$$

Coordinates of the intersection point are $(-0.588, 0.191)$.

<p>13 (i)</p>	$S = \frac{1}{2} H (2x + 8k) = \frac{1}{2} (2x + 8k) \sqrt{(4k)^2 - x^2}$ $= (x + 4k) \sqrt{16k^2 - x^2} \text{ (shown)}$ 	
<p>(ii)</p>	$\frac{dS}{dx} = \sqrt{16k^2 - x^2} - (x + 4k)x(16k^2 - x^2)^{-\frac{1}{2}}$ $= \frac{-2x^2 - 4kx + 16k^2}{\sqrt{16k^2 - x^2}} = 0$ $x^2 + 2kx - 8k^2 = 0$ $(x - 2k)(x + 4k) = 0$ $\Rightarrow x = 2k \text{ or } x = -4k \text{ (N.A., } x > 0)$ $\frac{dS}{dx} = \frac{-2x^2 - 4kx + 16k^2}{\sqrt{16k^2 - x^2}} = \frac{-2(x^2 + 2kx - 8k^2)}{\sqrt{16k^2 - x^2}}$ $\frac{d^2S}{dx^2} = -2 \left[\frac{\sqrt{16k^2 - x^2} (2x + 2k) - (x^2 + 2kx - 8k^2) \frac{1}{2} (16k^2 - x^2)^{-\frac{1}{2}} (-2x)}{16k^2 - x^2} \right]$ <p>when $x = 2k$</p> $\frac{d^2S}{dx^2} = -2 \frac{\sqrt{12k^2} 6k}{12k^2} = -\sqrt{12} < 0$ <p>Area of trapezium is maximum when $x = 2k$.</p>	
<p>(iii)</p>	<p>Using similar triangles:</p> $\frac{h}{H} = \frac{w - 2k}{2k}$ <p>When $x = 2k$, $H = \sqrt{12k^2} = 2\sqrt{3}k$</p> $w = \frac{2kh}{H} + 2k = \frac{h}{\sqrt{3}} + 2k$ $\therefore V = \frac{3}{2} h (4k + 2w) = \frac{3}{2} h \left(4k + 2 \left(\frac{h}{\sqrt{3}} + 2k \right) \right)$ $= 3h \left(4k + \frac{h}{\sqrt{3}} \right) \text{ (shown)}$ 	

<p>(iv)</p>	$\frac{dV}{dh} = \left(4k + \frac{h}{\sqrt{3}}\right)3 + 3h\left(\frac{1}{\sqrt{3}}\right)$ $= 12k + \frac{6h}{\sqrt{3}}$ <p>When $h = \sqrt{3}k$, $\frac{dV}{dh} = 18k$</p> $\frac{dh}{dt} = \frac{dh}{dV} \times \frac{dV}{dt} = \frac{0.2}{18k} = \frac{1}{90k} m/s$	
--------------------	---	--